

THE STATISTICAL MECHANICS OF BLACK HOLE THERMODYNAMICS *

Rafael D. Sorkin

Instituto de Ciencias Nucleares, UNAM, A. Postal 70-543, D.F. 04510, Mexico

and

Department of Physics, Syracuse University, Syracuse, NY 13244-1130, U.S.A.

internet address: sorkin@suhep.syr.edu

Abstract

Although we have convincing evidence that a black hole bears an entropy proportional to its surface (horizon) area, the “statistical mechanical” explanation of this entropy remains unknown. Two basic questions in this connection are: what is the microscopic origin of the entropy, and why does the law of entropy increase continue to hold when the horizon entropy is included? After a review of some of the difficulties in answering these questions, I propose an explanation of the law of entropy increase which comes near to a proof in the context of the “semiclassical” approximation, and which also provides a proof in full quantum gravity under the assumption that the latter fulfills certain natural expectations, like the existence of a conserved energy definable at infinity. This explanation seems to require a fundamental spacetime discreteness in order for the entropy to be consistently finite, and I recall briefly some of the ideas for what the discreteness might be. If such ideas are right, then our knowledge of the horizon entropy will allow us to “count the atoms of spacetime”.

When I first learned of the thermodynamics of black holes, and specifically of the fact that a black hole possesses an entropy proportional to its horizon area, my reaction (after thinking about it a while) was that this was just as if the horizon were divided into small “tiles” of a fixed size, with each tile carrying roughly one bit of information. To see that this would lead to the correct proportionality law, imagine a 0 or 1 engraved

* Published in the proceedings of the Chandra Symposium: “Black Holes and Relativistic Stars”, ed. R. M. Wald, held December, 1996, Chicago, (U. of Chicago Press), pp. 177-194.

on each tile. If there were A tiles, then the number of possible configurations would be $N = 2 \times 2 \times 2 \cdots \times 2 = 2^A$, and the corresponding entropy would be $S = \log N = A \log 2$ (taking Boltzmann's constant k equal to one). In order to get the numerical coefficient right, the tile size would have to be around 10^{-65}cm^2 , that is, it would have to be of order unity in units with $c = \hbar = 8\pi G = 1$.

Of course no one believes (or would admit to believing) that the black hole horizon is painted with tiny 0's and 1's, but the suggestion remains that, in some less artificial manner, a cutoff occurs at around the Planck scale, and the microscopic degrees of freedom proper to the horizon carry about one bit of information per horizon "atom". In order for such an explanation to work, it would have to provide convincing answers to two principal questions: what degrees of freedom does this information capture (to what N is one really referring when one says $S = k \log N$), and why does the total entropy still increase when black holes are involved (why does the Second Law of Thermodynamics continue to hold)?

The explanation that would resolve these twin questions is what I mean by the "statistical mechanics behind black hole thermodynamics". We can hope that in the process of arriving at such an explanation, we will learn something important about the nature of spacetime on small scales, just as the quest for the statistical mechanics of a box of gas taught us something important about the nature of ordinary matter on atomic scales, revealing the existence of atoms, their sizes, and something about their structure and quantum nature.

Equally, we may hope that the investigation of these questions will shed new light on statistical mechanics itself, and specifically on the meaning of entropy. For example, the experience so far has been that no derivation of the Second Law can even get started without first applying some form of coarse-graining to define the entropy, and the choice of coarse-graining seems to introduce an unwelcome element of subjectivity into the foundations of statistical mechanics. But a black hole affords an obvious objective way to coarse-grain, namely neglect whatever is inside the horizon. Or to take another example, it is the possibility of fluctuations in the entropy that distinguishes the statistical mechanical picture from the thermodynamical one, but in practice such fluctuations are ordinarily too small to be observable. With black holes, on the other hand, one can, at least in principle, arrange for arbitrarily large fluctuations to occur [1].

Before we address the statistical mechanics of black holes as such, let me remind you very briefly of their thermodynamic properties in a little more detail. In a process like stellar collapse, a black hole is said to form when a region of spacetime develops from which signals can no longer escape. Within the context of classical General Relativity the subsequent occurrence of a singularity is then inevitable, but it is expected to form inside the black hole, so that it is hidden from the view of distant astronomers (see the discussion in [2]).

The black hole’s *horizon* H is by definition the three dimensional surface separating the interior of the black hole from its exterior. Formally, we have $H = \partial(\text{past } \mathcal{I}^+)$, where \mathcal{I}^+ (called “future null infinity”) is defined as the set of ideal points at infinity at which outgoing light rays terminate. Thus the past of \mathcal{I}^+ consists of all events that can send light rays to infinity; and H is its boundary. From this definition alone certain mathematical facts follow, including the fact that H is a continuous surface (a C^0 manifold) which, although it may not be smooth (because of caustics), nevertheless is null almost everywhere in the sense that it is a union of null geodesics which never leave H as they propagate into the future. (These ruling geodesics are photon world lines that hover on the horizon, balanced precariously between escaping to infinity and being pulled into the singularity.) When we speak of the “area of the horizon”, we mean the area of the *cross-section* in which H intersects some spacelike (or possibly null) hypersurface Σ on which we seek to evaluate the entropy: *

$$A := \text{Area}(H \cap \Sigma).$$

Now an abundance of evidence indicates that there is associated with an event horizon of this sort an entropy of

$$S_{BH} = \frac{2\pi A}{l^2}, \tag{1}$$

where

$$l^2 = 8\pi G\hbar \tag{2}$$

* For this definition of A to make sense, it is necessary that $H \cap \Sigma$ not be too rough. I don’t know whether one can prove this for every possible black hole horizon H and smooth surface Σ , but “Geometric Measure Theory” guarantees at least that, for any given H , there exists a dense set of Σ ’s for which A is well defined [3].

is the square of the “rationalized Planck length”. The best known piece of evidence for this association is that the black hole radiates in precisely the right manner to be in equilibrium with a surrounding gas of thermal radiation at the temperature derived from (1) via the First Law of Thermodynamics, $dM = TdS$ (the Hawking radiation). But the evidence goes well beyond that, ranging from theorems in classical General Relativity that can be interpreted as the Zeroth through Third Laws of Thermodynamics applied to a black hole in the $\hbar \rightarrow 0$ limit, to computations directly yielding the entropy (1) in the “tree level” approximation of path-integral quantum gravity.

Many of these relationships carry over to other spacetime dimensions and other gravity theories than standard General Relativity, as emphasized in [4]. However, the evidence remains less complete and convincing in these other cases.* In particular one still lacks an analog for these other cases of the result that in standard General Relativity represents the $\hbar \rightarrow 0$ limit of the Second Law of Thermodynamics, namely the theorem that classically the total horizon area necessarily increases (or remains constant) as the hypersurface Σ on which it is evaluated moves forward in time. Because of the way that \hbar enters into the expressions (2) and (1), the black hole entropy goes to infinity in the classical limit, and so tends to dominate all other entropies. Therefore, this classical law of area increase can be interpreted as precisely what remains of the Second Law when \hbar tends to zero. (Unfortunately, the proof of area increase remains incomplete even in standard General Relativity, because it rests on the still unproven assumption of “cosmic censorship”.)

Notice that the law of area increase, even though it is valid only in the non-quantum limit, is a fully nonequilibrium result in the sense that no requirement of stationarity is imposed on the black holes or the classical matter with which they interact. In the complementary limit of fully quantum matter interacting with a near equilibrium black hole, there exist several arguments and thought experiments suggesting the impossibility of procuring a decrease in the total entropy (black hole plus entropy of surroundings) by any process in which the black hole evolves quasi-stationarily and remains essentially classical. (Later, I will propose a completely general proof of this impossibility.) Taken

* In a sense it is disappointing that black hole thermodynamics appears to be so robust: the less sensitive it is to the theoretical assumptions, the less capable of guiding us to the correct underlying theory of quantum gravity!

together, these results in special cases strongly suggest that the Second Law of Thermodynamics, $\Delta S \geq 0$, will continue to hold in general if we attribute to each black hole its corresponding entropy (1) and take for the total entropy, the sum of this horizon-area entropy with the entropy of whatever else may be present *outside* the black holes:

$$S_{tot} = S_{BH} + S_{outside} \quad \text{increases with time.} \quad (3)$$

Remark The above discussion has defined a black hole in causal terms, and correspondingly has taken the horizon to be what is more precisely referred to as the *event horizon*. A rather different concept is that of *apparent horizon*, whose definition generalizes the curvature properties of the Schwarzschild horizon, rather than its causal properties. In application to time-independent black holes, the two concepts coincide, but away from equilibrium they differ. It has been proposed [5] to identify the area that enters into the formula (1) as the area of the apparent horizon rather than the event horizon. The apparent horizon has the apparent advantage of being definable “quasi-locally”, whereas locating the event horizon requires in principle a knowledge of the entire future of the spacetime. On the other hand, it is precisely this “teleological” attribute of the event horizon that gives rise to the large entropy fluctuations alluded to above. Moreover, the concept of event horizon seems more fundamental than that of apparent horizon, and therefore more robust in relation to possible discrete replacements for spacetime such as the causal set [6]. In addition, the area of the apparent horizon will often jump discontinuously, which is not how one might expect entropy to behave. For these reasons, I will stick with the identification, horizon = event horizon, but it seems prudent to bear the other alternative in mind as well.

Now, how would we understand the nondecreasing character of S_{tot} if a black hole were an ordinary, nonrelativistic thermodynamic object? For example, suppose the black hole were really a warm brick or, say, a hot ball of hafnium. In that case (recalling the familiar plausibility arguments, as nicely presented in [4]), we would identify N_{BH} (the subscripts stand for “ball of hafnium”) as the number of internal micro-states of the ball, and $S_{BH} = \log N_{BH}$ as its logarithm. If we went on to assume that the ball was *weakly coupled* to its surroundings, then we could write (with N_{out} being the number of micro-states of the surroundings),

$$N_{tot} = N_{BH} \cdot N_{out} \quad (4a)$$

$$S_{tot} = S_{BH} + S_{out} \quad (4b)$$

(On the other hand, if the coupling was not weak, then the states of the two subsystems would not be able to vary independently, and so the estimate $N_{tot} = N_{BH} \cdot N_{out}$ would be inaccurate.) Under the further assumption that the dynamical evolution of the whole system was *ergodic* (exploring all of the available state-space) and *unitary* (so preserving state-space volume as measured by number of states), we would conclude that the time spent in any region of state-space was proportional to the number of states in that region:

$$dwell\ time \propto N_{tot}, \tag{5}$$

so that* the probability of a transition in which N_{tot} increased would be more likely than one in which it decreased, the discrepancy being overwhelming for large entropy differences because $\Delta S \gg 1 \Rightarrow e^{\Delta S} \gg \gg 1$.

So what goes wrong with this reasoning if we try to apply it to a black hole? In the first place, it is at least peculiar that the number of black hole states would be proportional to e^{Area} rather than e^{Volume} as for other thermodynamic systems. This peculiarity becomes more troubling if we consider the example of the Oppenheimer-Snyder spacetime, in which a Friedmann universe of *arbitrary size* is joined onto the interior of a Schwarzschild black hole of arbitrary mass. (See the remarks in [9].) The

* The plausibility argument here could be made stronger if we assumed a form of “detailed balance” for the effective Markov process operating on the macro-states, or better “meso-states” (cf. [7]). Then it would follow immediately that the probability (per unit time, say) of a transition from state 1 to state 2, given that the system was currently in state 1, would be greater by $e^{S_2 - S_1}$ than that of the reverse transition, given that the system was currently in state 2.

It is interesting that such a Markovian model of the evolution appears easier to justify in a classical framework, where the system really is in a definite (albeit unknown) micro-state at any instant. In the quantum case, in contrast, there is no reason to believe that the state-vector normally lies within any particular eigensubspace of a macroscopic observable like the temperature distribution. Thus, although the *definition* of entropy works better in the quantum case (because “number of micro-states” really makes sense there), the argument for its *increase* seems to proceed more happily in the classical setting. To recover a Markovian picture in the quantum case as well, one might have recourse to a sum-over-histories interpretation of the formalism, or to the closely related picture in which coarse-grained histories are obtained from sequences of projection operators as in [8].

existence of such solutions means that the number of possible *interior* states for a black hole is really infinite, which is certainly not consistent with any formula like $S = \log N$.

A second set of problems concerns *ergodicity* and *internal equilibrium*. The course of events inside a collapsing star leads classically to a singularity, and it is not at all obvious that this is consistent with an ergodic exploration of all available states, including for example the Oppenheimer-Snyder states just described. But even if quantum effects did restore ergodicity, there would remain a problem with equilibrium. Although I did not stress it above, an assumption of internal equilibrium (or partial equilibrium) is needed in order to deduce the entropy from the values of a few macroscopic variables. For example, knowing only the surface temperature of the ball of hafnium would not at all allow you to deduce its entropy unless you added the assumption of internal equilibrium; if this assumption were mistaken, you might find the apparent entropy suddenly starting to decrease, because the interior of the ball was much colder than you had assumed. In the same way, mere knowledge of the external appearance of a black hole tells you little about its interior, and since realistic black holes would seem to be far from internal equilibrium, this is another reason to doubt whether the type of state-counting utilized in (4) and (5) can carry over to black holes with N_{BH} interpreted as the number of interior states.

A related problem with such state-counting is that our earlier assumption of *weak coupling* between the subsystem and its environment is doubly wrong in the case of a black hole. The coupling from outside to inside is not weak but very strong, while the reverse coupling is not so much weak as nonexistent! Indeed this last observation points up the fact that conditions in the interior should be irrelevant, almost by definition, to what goes on outside. And since the Second Law, as ordinarily formulated for black holes, makes no reference to conditions inside, it seems especially strange that it should have anything to do with counting interior states. (In contrast, the temperature distribution inside our ball of hafnium can make a big difference in its interaction with the outside world, as we have seen, and it is impossible to specify its entropy without saying something about the internal conditions, if not explicitly, then implicitly via the assumption of internal equilibrium.)

Finally there is the vexed question of *unitarity*. Many people refer to this in terms of an “information puzzle”, the puzzle being that the facts apparently belie their belief that there must be a well-defined unitary S-matrix for the *exterior* region alone; but for our

purposes here, the existence of such an S-matrix is irrelevant, because all we are interested in is the Second Law, and that certainly imposes no such requirement.* Rather, what the ordinary statistical mechanical reasoning requires is *overall* unitarity (for exterior and interior regions together), and the difficulty lies in combining this overall unitarity with the mode of explanation that would locate the entropy of the black hole in the multiplicity of its interior micro-states. If the latter were correct, then a contradiction with unitarity would arise in the process of evaporation of a black hole by Hawking radiation. In fact, since the black hole loses entropy as it shrinks (its surface area decreases), its number of internal states would have to go down as well, and at some point there would no longer be enough of them to support the correlations with the radiated particles which are needed if the *overall* evolution is to remain unitary.

At least this conclusion is inevitable if we accept the semiclassical description of the radiation as consisting of correlated pairs A and \bar{A} , the first of which goes off to infinity while the second falls into the singularity. In this approximation of quantum field theory on a fixed, background black hole metric, the emitted particles A taken alone are described by a highly impure state of the exterior field, and the unitarity of the overall quantum evolution is restored only when the $A \leftrightarrow \bar{A}$ correlations are taken into account. (One sometimes says that the particles A are “entangled” with the particles \bar{A} .)

In order for these entanglement correlations to exist, however, it is necessary that the interior of the black hole support a quantum state-space of dimension at least $e^{S_{rad}}$, where S_{rad} is the entropy of the emitted thermal radiation. If the number of internal states really diminished as the black hole shrank then this would cease to be possible, and so unitarity could be maintained only if the $A \leftrightarrow \bar{A}$ correlations were transferred to ones of the form $A \leftrightarrow B$ between *emitted* particles. Ultimately, if the black hole were allowed to evaporate fully, then all of the $A \leftrightarrow \bar{A}$ correlations would have to be transferred to the outside; and thus unitary evolution of the whole system, *together with* the assumption that black hole entropy counts the number of internal states, would require the external evolution *alone* to be unitary, at least in the S -matrix sense.

* On the contrary, a unitary evolution of the exterior region would be inimical to the type of explanation of the Second Law I will propose below.

Now, it is possible to imagine mechanisms by which the required $A \leftrightarrow B$ correlations could arise, but it is much less easy to imagine how the $A \leftrightarrow \bar{A}$ correlations could disappear, since they arise in an approximation which should remain good in full quantum gravity. But if they don't disappear then the quantum state would have to stay "entangled", which in turn would require the number of internal states to remain much larger than allowed by the formula (1). Hence one has either to abandon the interpretation of black hole entropy in terms of internal states or to grant that the overall quantum evolution of the system, black hole + environment, is nonunitary (or both).

Despite the seeming inevitability of this conclusion, many workers still hope to evade it, but in seeking to do so, they are driven to rather desperate maneuvers, such as hypothesizing a new "complementarity" according to which the internal state-space would be of a different dimensionality for observers outside the black hole than for observers inside of it. Such conceptual difficulties notwithstanding, the attempt to imagine how the external region alone can possess a unitary S -matrix (or perhaps even a dynamics which remains unitary during all intermediate stages) has led to some interesting suggestions, including the suggestion [10] that at very small distances the only variables that remain are purely geometrical ones, with all other distinctions (color, flavor, generation, etc.) being washed out.

By far the greatest effort toward providing a unitary statistical mechanics for black hole thermodynamics has been exerted within the context of string theory (which is not to say that strings are necessarily tied to a unitary explanation, any more than any other approach is, cf. [11]). In fact most of the effort has been directed to only one of the two questions I emphasized at the outset, namely the question of what degrees of freedom one must count in order to obtain the formula (1). The earliest calculation of this sort that I know of was performed by Steve Carlip [12] for a black hole in 2+1 dimensional gravity. Although this calculation did not actually use a stringy version of gravity, it used string technology to count certain "gauge" degrees of freedom defined on the horizon, obtaining (1) with precisely the thermodynamically required coefficient. This would tend to agree with the suggestion that the entropy is localized at the horizon, rather than inside the black hole. More recently, some calculations in string theory proper have also obtained equation (1) for zero temperature black holes in certain higher dimensional theories of gravity coupled to special combinations of gauge fields chosen to make supersymmetric solutions possible. These more recent calculations do not actually work with black holes,

rather they count certain “membrane” states in a flat-space limit and obtain a formula which can be interpreted as the analytic continuation of (1) to that limit. (See [13] for a review of this work.)

To the extent that the physical picture behind these stringy calculations can be extrapolated from flat spacetime to genuine black hole geometries (which assumes in particular that the “phase-transition” accompanying the formation of the horizon would not interfere with analyticity), they suggest that the degrees of freedom contributing to the entropy are associated with certain types of membranes. In contrast to Carlip’s calculation, they also could be taken to suggest that the relevant degrees of freedom are those of the black hole as a whole (as opposed to being localized at the horizon), but since there are no horizons in flat space, such a suggestion would have to be very tentative at best. Beyond this, the stringy calculations do not (as far as I can see) shed much light on the question that to me is most central about the entropy of black holes: why is it finite at all? We will see later that certain identifiable contributions to the entropy that arguably must be present are infinite in the absence of a short-distance cutoff at the horizon. One might ask whether these contributions are present in the string theory picture, and if so, how the requisite cutoff arises.

Concerning the derivation of the Second Law and the objections raised above, string theory apparently has little to say. If, at the end of the day, it is able to produce a unitary dynamics underlying quantum gravity, and if it is able to count the states of a black hole in this framework (and also explain why a weak coupling, equilibrium approximation is really valid after all, etc.), then the derivation of the Second Law will be reduced to the discussion one can find in any thoughtful textbook on statistical mechanics. However, we have seen that the demand for unitarity in particular, seems to drive one to a new kind of inside–outside complementarity that so far no one knows how to formulate. For this reason it seems impossible at this stage to address the question of the Second Law within the confines of any unitary theory that identifies the black hole entropy with the states of the black hole taken as a whole. Let us return, then, to our two main questions — “Why does S increase?” and “What does S count?” — and consider them in the order just given, hoping that an answer to the first will suggest an answer to the second.

Why does S increase?

What I want to describe now is an old proposal [14] [15] that would derive the Second Law by appealing directly to the property of black holes that makes them *different* from all other objects, the fact that what goes on outside the horizon takes place without any reference to what goes on inside. Classically this independence is exact, and quantum mechanically it can still be expected to hold to an excellent approximation, despite some quantum blurring of the horizon. But if the exterior region really has a well-defined autonomous dynamics, then it is particularly natural to “coarse-grain away” the interior region and seek to identify the entropy that enters into the Second Law with the entropy of some effective quantum density-operator describing the exterior situation.*

More specifically, what we would like to do is evaluate the entropy on a hypersurface Σ like that discussed earlier, but with the difference that our new Σ terminates where it meets the horizon. We would like to associate an effective density-operator $\hat{\rho}(\Sigma)$ with each such hypersurface and to prove that

$$S(\Sigma'') \geq S(\Sigma')$$

whenever Σ'' lies wholly to the future of Σ' . We will see, in fact, that this approach to the question leads to two proofs of entropy increase, a more fully worked out one which applies in the semiclassical approximation [15] (cf. the remarks in [16] and the related remarks in [17] and [18]), and a more sketchy one which applies in full quantum gravity [14] [15]. It also leads to the conclusions that the entropy is localized (to the extent that entropy can ever be localized) at or just outside the horizon [14] [19] [16], and that S owes its finiteness to a fundamental spacetime discreteness or “atomicity” [14] [19].

Let us take first the case of a quasi-classical, quasi-stationary black hole, by which I mean that we ignore quantum fluctuations in M and the other black hole parameters

* This is not meant to imply that on philosophical grounds, we must neglect the interior region because “we” cannot observe it. Rather my attitude would be that any coarse-graining at all is valid if it leads to a useful definition of entropy. The horizon is nevertheless special in this connection because, as we will see, its special causal properties make it possible to draw conclusions that could not be drawn with other types of coarse-graining.

and we assume that the spacetime geometry can be well approximated at any stage by a strictly stationary metric. [In other words, we work in the framework of “quantum field theory in curved spacetime”. Notice that the requirement of approximate stationarity applies only to the metric; the matter-fields (among which we may include gravitons) can be doing anything they like.] In such a situation there exists for the matter fields outside the black hole a unique thermal (or “KMS”) state $\hat{\rho}^0$, known as the “Hartle-Hawking state” and expressible as

$$\hat{\rho}^0 \propto e^{-\beta \hat{E}}, \quad (6)$$

where \hat{E} represents the energy operator for the matter fields in the external region and β is the reciprocal of the thermodynamic temperature of the black hole.* (For simplicity, we may as well work with a nonrotating, charge-free black hole. Alternatively, one could just reinterpret \hat{E} as $\hat{E} - \omega \hat{J} - \phi \hat{Q}$.)

Now let us limit our consideration of the entropy to the hypersurfaces $\Sigma(t)$ of a foliation of the exterior region which is compatible with the Killing vector field ξ^a that expresses the stationarity of the metric. In other words, the $\Sigma(t)$ are the $t = \text{constant}$ surfaces of some coordinate system for which the metric is explicitly time-independent. (Notice, however, that it would *not* be appropriate to take t to be the Schwarzschild time coordinate, for example, since then the cross-section $\Sigma(t) \cap H$ would not move forward along the horizon as t increased.) To each hypersurface $\Sigma(t)$ there corresponds a quantum Hilbert space in which the field operators residing on $\Sigma(t)$ are represented (in particular the operator $\hat{E} = \hat{E}(t)$ of eq. (6)), and in which the quantum state $\hat{\rho}(t)$ can therefore be represented as a density operator. By identifying appropriately the hypersurfaces $\Sigma(t)$ with one another (the natural identification here being that induced by the Killing vector ξ^a), we identify their attached Hilbert spaces, and the dynamical change of $\hat{\rho}(t)$ with time becomes thereby a motion of $\hat{\rho}(t)$ within a single Hilbert space. (Notice that this evolution of $\hat{\rho}$ is well defined because the boundary H at which we have truncated the hypersurfaces Σ is an *event horizon*, across which no information can propagate. This

* This expression is inevitably not rigorous, if only because the theory of (interacting) quantum fields is itself not rigorous. A related comment is that, in a nonstationary background, there would be difficulties with the definition of \hat{E} , even in the case of a free field, as Adam Helfer pointed out to me after my lecture [20].

property of autonomous evolution would be lost if, say, H were replaced by a timelike surface, or if we tried to evolve $\hat{\rho}(t)$ *backward* in time.)

Now the key feature of the state (6) for us is that it is time-independent—as every equilibrium state must be by definition. What makes this so important is that given any well-defined (*but not necessarily unitary*) evolution law for density-operators $\hat{\rho}(t)$, and given any state $\hat{\rho}^0$ that is stationary with respect to this evolution, we can find a function $f(\hat{\rho}(t))$, defined for *arbitrary* states $\hat{\rho}(t)$, which is nondecreasing with time. (More precisely, this is true classically for an arbitrary Markov process, and true quantum mechanically under a certain auxiliary technical assumption.) In effect, every state “wants to evolve toward the stationary one”, and f is a kind of Lyapunov functional measuring how close $\hat{\rho}(t)$ has come to $\hat{\rho}^0$. Now, when the stationary state has the Gibbsian form (6), the quantity $f(\hat{\rho})$ turns out to be (up to an additive constant)

$$S(\hat{\rho}) - \beta \langle \hat{E} \rangle = \text{Tr } \hat{\rho} (\log \hat{\rho}^{-1} - \beta \hat{E}); \quad (7)$$

that is, it turns out to be the free energy up to a factor of $-1/T$.

The proof of this little known result is so remarkably simple (at least classically) that I cannot resist presenting it here, in the important special case where $\beta = 0$. To begin with, note that the function $f(x) \equiv x \ln x^{-1}$ is concave downward since $f''(x) = -1/x < 0$ in the range $0 \leq x \leq 1$. Now consider a Markov process whose probability of being in the k^{th} state at some moment of time is p_k . The quantity (7), when $\beta = 0$, is nothing but the entropy

$$S(p) = \sum_k p_k \log p_k^{-1} = \sum_k f(p_k),$$

so what we have to prove is that $S(p)$ increases when the p_k get replaced at some later time by $\sum_l T_{kl} p_l$, T_{kl} being the matrix of transition probabilities. But we have

$$\begin{aligned} S(p) &= \sum_k f(p_k) \\ &\rightarrow \sum_k f\left(\sum_l T_{kl} p_l\right) \\ &\geq \sum_{kl} T_{kl} f(p_l) \\ &= \sum_l f(p_l), \end{aligned}$$

where to get the inequality, we used the concavity of f together with the fact that

$$\sum_l T_{kl} = 1$$

because T preserves (when $\beta = 0$) the totally random state, $p_k \equiv 1$; and in the last step we used that

$$\sum_k T_{kl} = 1$$

(conservation of probability). When the stationary state p_k^0 is not uniform, the proof is almost as simple, only S gets replaced by

$$\sum_k p_k^0 f(p_k/p_k^0),$$

which for a thermal $p_k^0 = e^{-\beta E_k}$ is just

$$\sum_k p_k^0 \frac{p_k}{p_k^0} \log \frac{p_k^0}{p_k} = \sum_k p_k (-\beta E_k - \ln p_k) = S - \beta \langle E \rangle,$$

which is indeed the classical form of (7).

Returning to the black hole situation, it is now simple to see that the nondecreasing character of (7) entails that of the total entropy in our semiclassical, quasistationary approximation. To that end, consider the small change of state that occurs as the hypersurface Σ' moves slightly forward in time to Σ'' , and write now S_{out} for the entropy $S(\hat{\rho})$ of the exterior matter, in order to distinguish it from the entropy S_{BH} of the black hole itself. For the latter we have (from the ‘‘First Law of black hole thermodynamics’’) that

$$dS_{BH} = \beta dM, \tag{8}$$

M being the mass of the hole. As part of the semiclassical approximation, we also assume that the mass of the black hole adjusts itself in response to the quantum mechanical

expectation value of the energy it exchanges with the matter fields. Therefore (changing ‘ \hat{E} ’ to ‘ \hat{E}_{out} ’ for notational consistency) we have*

$$\langle \hat{E}_{out} \rangle + M = \text{constant}, \quad (9)$$

whence $dM = -d \langle \hat{E}_{out} \rangle$ and $dS_{BH} = -\beta d \langle \hat{E}_{out} \rangle$. Putting this together with the fact that the expression (7) must not decrease in the process (and observing that β in that expression represents a *fixed* parameter of the stationary metric), we can write for the infinitesimal change $\Sigma' \rightarrow \Sigma''$,

$$\begin{aligned} d(S_{out} - \beta \langle \hat{E}_{out} \rangle) &\geq 0 \\ dS_{out} - \beta d \langle \hat{E}_{out} \rangle &\geq 0 \\ dS_{out} + dS_{BH} &\geq 0 \\ d(S_{out} + S_{BH}) &\geq 0 \end{aligned} \quad (10)$$

Thus we have proved the Second Law in the semiclassical approximation, for arbitrary processes in which the black hole geometry changes sufficiently slowly. This class of situations includes all that I know of for which thought experiments have been done to check the Second Law (see [4] for some of them); however it does not cover black holes that are far from equilibrium, for example ones just formed by the coalescence of two neutron stars and still in the process of settling down to a stationary state. Notice also that in this proof we are not *deriving* the value of the horizon entropy, but only showing that the Second Law holds *if* we use the value (1) provided by thermodynamic arguments.

Finally, it should be added that the matter entropy $S(\hat{\rho})$ we have been working with is actually infinite, due to the entanglement between values of the quantum fields just

* One might worry that the two terms in this equation refer to two different energies, M to the total energy as defined at infinity, and $\langle \hat{E} \rangle$ to an energy defined with respect to the stationary background geometry. That this is not really a problem can be seen, for example, by imagining an infinitesimal mass m falling into the black hole from infinity. On one hand, this augments the mass M of the black hole by m ; on the other hand, by conservation of the conserved energy current $T_b^a \xi^b$ in the stationary background, the exterior energy E_{out} decreases by the same amount m as the mass passes through the horizon. A more direct proof that (classically) $dS_{BH} = -\beta dE_{out}$ may be extracted from the derivation following eq. (13) in the first reference of [21].

inside and just outside the horizon. In a little while we will consider this near-horizon entanglement entropy as a possible source of the black hole entropy itself, but for now it is just a nuisance since the divergence in S_{out} means that (7) diverges as well. Thus, making our proof rigorous would require showing that *changes* in (7) are nevertheless well-defined and conform to the temporal monotonicity we derived for that quantity. This probably could be done by introducing a high-frequency cutoff on the Hilbert space (using as high a frequency as needed in any given situation) and showing that the evolution of $\hat{\rho}$ remained unaffected because the high-frequency modes remained unexcited.*

To what extent can we expect to generalize these considerations to the case of greatest interest, that of a fully dynamical, quantum black hole (or holes)? In one respect, the situation actually simplifies, because the technical hypothesis of complete positivity is no longer needed. The complication, of course, is that the continued validity of some of our other assumptions will depend on the structure of the quantum gravity theory in which the proof is to be realized. Without knowing that structure in advance we can do no better than to list the features that would be needed in order for the proof to go through. In writing down this list, I will assume that the hypersurfaces Σ to which the entropy is being referred can be specified by some generally covariant prescription that continues to make sense in quantum gravity, at least in some sufficiently great subset of the situations for which we would want to formulate a Second Law. (For example, we might take advantage of the fact that a “box” is needed in order for thermodynamic equilibrium to be possible, and specify Σ as the boundary of the future of a freely chosen cross-section of the box, regarded as a timelike “world tube” of fixed geometry. [More precisely, Σ would be that part of the boundary lying inside of the box and outside of any future horizons

* Alternatively, perhaps one could show that the additive constant, $-\log \text{Tr } e^{-\beta H}$, omitted from (7) canceled the divergence in a well-defined sense. In order to make the proof rigorous, one would also have, for example, to specify an observable algebra for the exterior fields and a representation of that algebra in which the operators $\hat{\rho}$ and \hat{E} were well-defined (which in particular might raise the issue of boundary conditions near the horizon). In addition, one would have to prove that the autonomous evolution of $\hat{\rho}$ was “completely positive”, this being a condition for the applicability of the theorem that guarantees the nondecreasing nature of (7) in the quantum case. This probably could be done by expressing the autonomous evolution as a unitary transformation followed by a tracing out of interior degrees of freedom. See [15] for references and some further discussion of these points.

that were present.] Such a Σ would be “achronal”, though not strictly spacelike, and the temporal relationship of two Σ ’s specified in this way would follow directly from that of their cross-sections.) With respect to some such family of hypersurfaces the needed assumptions are these:

- there is defined a Hilbert space \mathcal{H} , and for each surface Σ , an effective density-operator $\hat{\rho}(\Sigma)$ acting in \mathcal{H} ,
- $\hat{\rho}$ evolves autonomously in the sense that $\hat{\rho}(\Sigma'')$ is determined by $\hat{\rho}(\Sigma')$ whenever Σ'' lies to the future of Σ' ,
- there exists an operator \hat{E} defined at the boundary of the system (or in any case without direct reference to the region inside the black hole) and yielding the total conserved energy,
- the operator $\hat{\rho}_{E_0} = \theta(E_0 - \hat{E})$ is preserved by the autonomous evolution,
- $(\forall E_0) \dim(\mathcal{H}_{\hat{E} < E_0}) < \infty$, where $\mathcal{H}_{\hat{E} < E_0}$ is the subspace of \mathcal{H} with total energy less than E_0 .

Depending on your expectations, hopes or fears for quantum gravity, you may find different ones of these assumptions more or less plausible or palatable. For me all are at least plausible, albeit the entire framework of Hilbert space and operator observables is unlikely to be reproduced in an exact form in quantum gravity. In any case, if we accept these assumptions, then it follows easily that the quantity

$$S_{tot} \equiv \text{Tr} \hat{\rho} \log \hat{\rho}^{-1} \tag{11}$$

is a nondecreasing function of Σ . (The underlying mathematical theorem is proved in [22] and also in [15], where some further discussion of the above assumptions may be found as well.) This, then, will establish the Second Law (3) if we can show in addition that S_{tot} is a sum of two terms, one associated with the horizon and one with the exterior region.

Why is S a sum?

So why would $S_{tot} = \text{Tr} \hat{\rho} \log \hat{\rho}^{-1}$ take the form of the sum $A/l^2 + S_{surroundings}$ when $\hat{\rho}$ describes the exterior state including all fields, both gravitational and nongravitational?

For this to occur, there would need to be a great many degrees of freedom just outside or “contained in” the horizon (one might expect it to be thickened due to quantum effects), making their own identifiable contribution to S_{tot} . And of course this contribution would have to be proportional to the horizon area in order to agree with equation (1). Three ideas that have been proposed for what the horizon degrees of freedom might be are:

- (a) the geometrical shape of the horizon itself,
- (b) the modes of quantum fields propagating just outside the horizon,
- (c) the fundamental degrees of freedom of the substratum,

where by “substratum”, I mean the underlying structure(s) of whatever turns out to be the true theory of quantum gravity. Of course these possibilities are not necessarily exclusive of each other. All three types of contribution might be present, and they might also overlap significantly (for example, substratum degrees of freedom might show up in an effective description as geometrical variables describing the horizon shape). In concluding this article, I would like to say a few words about each of these possibilities and how they bear on the question of spacetime discreteness.

An attractive feature of the first proposal [14] [23] is that it offers a geometrical explanation for the very geometrical relationship (1). Unfortunately, it is not easy to estimate the magnitude of the quantum fluctuations in the horizon shape, but there are indications [21] that they become significant, not at the Planck scale l (as one might have expected), but at the much larger scale $\lambda_0 \sim (Ml^2)^{1/3}$. Such fluctuations would in effect spread the horizon into a shell of thickness λ_0 harboring a potentially unlimited source of entropy. In fact, the density of fluctuation modes diverges as their transverse wave-number goes to infinity. This means that, without any cutoff, the entropy residing in the shape degrees of freedom would presumably be infinite. On the other hand, the very rapidity of the modes’ growth rate also means that most of the modes with wavelength greater than any specified lower bound λ_{min} have $\lambda \sim \lambda_{min}$. In consequence, a result like (1), (2) emerges automatically (at least in crudely estimated order of magnitude) if one does assume a cutoff of magnitude $\lambda_{min} \sim l$.

The second proposal leads to very similar conclusions. Here, if we work in the approximation of a fixed background geometry, and if we limit ourselves to free quantum fields, then we can actually compute the entropy $S_{out} = - \text{Tr } \hat{\rho} \log \hat{\rho}$ of the field, and we

obtain [14] [23] the result $S \sim A/\lambda_{min}^2$ where λ_{min} is the cutoff. In this case, we recognize the area law explicitly, and we see again that we must choose $\lambda_{min} \sim l$ in order to recover an entropy of the correct order of magnitude.

Unlike for proposal (a), where the effect of shape fluctuations on the density operator $\hat{\rho}(\Sigma)$ is not completely clear, a contribution of type (b) must necessarily be present in the entropy (11). In conjunction with the type (a) contribution, this leads to an argument for the inevitability of spacetime discreteness: under the assumption of a true continuum, the type (b) entropy could fail to be infinite only if it provided its own cutoff by inducing, and thereby coupling to, horizon fluctuations of type (a), but then these would take over and provide an entropy that therefore would still be infinite. *

Another facet of proposal (b) which should be mentioned here is the apparent difficulty that, since each field makes its own contribution, the black hole entropy would seem to depend on the number of species of fundamental fields in nature. However, this conclusion is inevitable only if we ignore the coupling of field fluctuations to horizon shape and also hold fixed the value of the cutoff λ_{min} . But at fixed cutoff, not only S but also the Planck length (2) will be affected by the addition of a new species, and it appears [25] that this dependence is just what is needed to maintain the relationship (1) unchanged.

Of course the need for a cutoff bespeaks an underlying spacetime discreteness, and so, to my mind, the most intriguing possibilities of type (c) are those in which the substratum has a discrete character. In such a theory one could hope to derive the entropy, not just by counting discrete quantum states of physically continuous variables, but by counting certain discrete physical elements themselves (compare the fact that the entropy of a box of gas is, up to a logarithmic factor, just the number of molecules it contains). For example, in causal set theory, one might count the number of causal links crossing the horizon (near Σ), or in canonical quantum gravity in the loop representation, one might count the number of loops cut by the horizon (within Σ). In any such case, the result would be something like the horizon area in units set by the fundamental discreteness scale. Thus, to reduce the evaluation of the entropy (1) to a counting exercise of this sort would be to open up a direct path to learning the value of the fundamental length.

* For a partly different interpretation see [24].

And if that were to occur, then the quest for the statistical mechanics of black hole thermodynamics would certainly have led us to something of interest: it would have led us to the atoms of spacetime itself.

This research was partly supported by NSF grant PHY-9600620.

References

- [1] R.D. Sorkin and D. Sudarsky (in preparation).
- [2] R. Penrose, contribution to this volume.
- [3] Frank Morgan, *Geometric Measure Theory: A Beginner's Guide*, second edition (Academic Press 1988), Section 4.11. I thank G.J. Galloway for this information.
- [4] R.M. Wald, contribution to this volume, and references therein; "Entropy and Black Hole Thermodynamics", *Phys. Rev. D***20**: 1271-1282 (1979).
- [5] Sean A. Hayward, "General laws of black-hole dynamics", *Phys. Rev. D* **49**:6467-6474 (1994).
- [6] L. Bombelli, J. Lee, D. Meyer and R.D. Sorkin, "Spacetime as a Causal Set", *Phys. Rev. Lett.* **59**:521-524 (1987).
- [7] R.D. Sorkin, "Stochastic Evolution on a Manifold of States", *Ann. Phys. (New York)* **168**: 119-147 (1986).
- [8] C.J. Isham, "Quantum Logic and the Histories Approach to Quantum Theory", *J. Math. Phys.* **35**: 2157-2185 (1994) <e-print archive: gr-qc/9308006>.
- [9] R.D. Sorkin, R.M. Wald and Zhang Zh.-J., "Entropy of Self-Gravitating Radiation", *Gen. Rel. Grav.* **13**:1127-1146 (1981).
- [10] G. 't Hooft, "The Black Hole Interpretation of String Theory", *Nuc. Phys. B***335**: 138-154 (1990).
- [11] S.W. Hawking, contribution to this volume.
- [12] S. Carlip, "Statistical Mechanics of the (2+1)-dimensional Black Hole", *Phys. Rev. D* **51**: 632-637 (1995) <e-print archive: gr-qc/9409052>.
- [13] G. Horowitz, contribution to this volume.
- [14] R.D. Sorkin, "On the Entropy of the Vacuum Outside a Horizon", in B. Bertotti, F. de Felice and A. Pascolini (eds.), *Tenth International Conference on General Relativity*

- and Gravitation (held Padova, 4-9 July, 1983), Contributed Papers*, vol. II, pp. 734-736 (Roma, Consiglio Nazionale Delle Ricerche, 1983).
- [15] R.D. Sorkin, “Toward an Explanation of Entropy Increase in the Presence of Quantum Black Holes”, *Phys. Rev. Lett.* **56**, 1885-1888 (1986).
- [16] W.H. Zurek and K.S. Thorne, “Statistical Mechanical Origin of the Entropy of a Rotating, Charged Black Hole”, *Phys. Rev. Lett.* **54**: 2171-2175 (1985).
- [17] R.D. Sorkin, “On the Meaning of the Canonical Ensemble”, *Int. J. Theor. Phys.* **18**:309-321 (1979).
- [18] R.D. Sorkin, “A Simplified Derivation of Stimulated Emission by Black Holes”, *Classical and Quantum Gravity* **4**, L149-L155 (1987).
- [19] G. 't Hooft, “On the quantum structure of a black hole”, *Nuclear Phys. B* **256**:727-745 (1985).
- [20] A.D. Helfer, “The Stress-energy Operator”, *Class. Quant. Grav.* **13**: L129-L134 (1996).
- [21] R.D. Sorkin, “Two Topics concerning Black Holes: Extremality of the Energy, Fractality of the Horizon”, in S.A. Fulling (ed.), *Proceedings of the Conference on Heat Kernel Techniques and Quantum Gravity, held Winnipeg, Canada, August, 1994*, pp. 387-407 (Discourses in Mathematics and its Applications, #4) (University of Texas Press, 1995) <e-print archive: gr-qc/9508002>; and
 “How Wrinkled is the Surface of a Black Hole?”, in David Wiltshire (ed.), *Proceedings of the First Australasian Conference on General Relativity and Gravitation*, held February 1996, Adelaide, Australia, pp. 163-174 (University of Adelaide, 1996) <e-Print archive: gr-qc/9701056>;
 A. Casher, F. Englert, N. Itshaki, S. Massar and R. Parenti, “Black Hole Horizon Fluctuations” <e-Print archive: hep-th/9606106>.
- [22] Woo Ching-Hung, “Linear Stochastic Motions of Physical Systems”, Berkeley University Preprint, UCRL-10431 (1962).
- [23] L. Bombelli, R.K. Koul, Lee J. and R.D. Sorkin, “A Quantum Source of Entropy for Black Holes”, *Phys. Rev. D* **34**, 373-383 (1986).
- [24] J.D. Bekenstein, “Do we understand black hole entropy?”, in *The Seventh Marcel Grossmann Meeting on Recent Developments in Theoretical and Experimental General Relativity, Gravitation and Relativistic Field Theories*, proceedings of the MG7 meeting, held Stanford, July 24–30, 1994, edited by R.T. Jantzen, G. Mac Keiser and R. Ruffini (World Scientific 1996) <e-print archive: gr-qc/9409015>.
- [25] L. Susskind and J. Uglum, “Black hole entropy in canonical quantum gravity and superstring theory”, *Phys. Rev. D* **50**:2700-2711 (1994) <e-print archive: hep-th/9401070>;
 T. Jacobson, “Black Hole Entropy and Induced Gravity”, < gr-qc/9404039 >.